

订购热线: 4008-898-798

Anti-F3 antibody

Cat. No. ml262373

Package 25 μl/100 μl/200 μl

Storage -20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Product overview

Description Anti-F3 rabbit polyclonal antibody

Applications ELISA, IHC

Immunogen Synthetic peptide of human F3

ReactivityHumanContent1.1 mg/mlHost speciesRabbit

Ig classImmunogen-specific rabbit IgGPurificationAntigen affinity purification

Target information

Symbol F3

Full name coagulation factor III, tissue factor

Synonyms TF; TFA; CD142

Swissprot P13726

Target Background

This gene encodes coagulation factor III which is a cell surface glycoprotein. This factor enables cells to initiate the blood coagulation cascades, and it functions as the high-affinity receptor for the coagulation factor VII. The resulting complex provides a catalytic event that is responsible for initiation of the coagulation protease cascades by specific limited proteolysis. Unlike the other cofactors of these protease cascades, which circulate as nonfunctional precursors, this factor is a potent initiator that is fully functional when expressed on cell surfaces. There are 3 distinct domains of this factor: extracellular, transmembrane, and cytoplasmic. This protein is the only one in the coagulation pathway for which a congenital deficiency has not been described. Alternate splicing results in multiple transcript variants.




订购热线: 4008-898-798


Applications

Immunohistochemistry

Predicted cell location: Cytoplasm and Cell membrane

Positive control: Human tonsil Recommended dilution: 40-250





The image on the left is immunohistochemistry of paraffin-embedded Human tonsil tissue using ml262373(F3 Antibody) at dilution 1/60, on the right is treated with synthetic peptide. (Original magnification: ×200)

ELISA

Recommended dilution: 5000-10000


联系电话: 4008-898-798, 021-61725725


联系QQ: 2881505695, 2881505696

邮箱: mlbio_cn@yeah.net 网址: www.mlbio.cn

Predicted cell location: Cytoplasm and Cell membrane

Positive control: Human breast cancer Recommended dilution: 40-250





The image on the left is immunohistochemistry of paraffin-embedded Human breast cancer tissue using ml262373(F3 Antibody) at dilution 1/60, on the right is treated with synthetic peptide. (Original magnification: ×200)