

订购热线: 4008-898-798

Anti-HDAC5 antibody

Cat. No. ml262793

Package 25 μ l/100 μ l/200 μ l

Storage -20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Product overview

Description Anti-HDAC5 rabbit polyclonal antibody

Applications ELISA, IHC

Immunogen Synthetic peptide of human HDAC5

ReactivityHuman, MouseContent0.9 mg/mlHost speciesRabbit

Ig classImmunogen-specific rabbit IgGPurificationAntigen affinity purification

Target information

Symbol HDAC5

Full name histone deacetylase 5

Synonyms HD5; NY-CO-9
Swissprot Q9UQL6

Target Background

Histones play a critical role in transcriptional regulation, cell cycle progression, and developmental events. Histone acetylation/deacetylation alters chromosome structure and affects transcription factor access to DNA. The protein encoded by this gene belongs to the class II histone deacetylase/acuc/apha family. It possesses histone deacetylase activity and represses transcription when tethered to a promoter. It coimmunoprecipitates only with HDAC3 family member and might form multicomplex proteins. It also interacts with myocyte enhancer factor-2 (MEF2) proteins, resulting in repression of MEF2-dependent genes. This gene is thought to be associated with colon cancer. Two transcript variants encoding different isoforms have been found for this gene.



订购热线: 4008-898-798


Applications


Immunohistochemistry

Predicted cell location: Cytoplasm and Nucleus

Positive control: Human gastric cancer

Recommended dilution: 25-100





The image on the left is immunohistochemistry of paraffin-embedded Human gastric cancer tissue using ml262793(HDAC5 Antibody) at dilution 1/20, on the right is treated with synthetic peptide. (Original magnification: ×200)

ELISA

Recommended dilution: 5000-10000


联系电话: 4008-898-798, 021-61725725


联系QQ: 2881505695, 2881505696

邮箱: mlbio_cn@yeah.net 网址: www.mlbio.cn

Predicted cell location: Cytoplasm and Nucleus

Positive control: Human tonsil Recommended dilution: 25-100





The image on the left is immunohistochemistry of paraffin-embedded Human tonsil tissue using ml262793(HDAC5 Antibody) at dilution 1/20, on the right is treated with synthetic peptide. (Original

magnification: ×200)