

Anti-HMGCL antibody

Cat. No.	ml122488
Package	25 µl/100 µl/200 µl
Storage	-20°C, pH7.4 PBS, 0.05% NaN ₃ , 40% Glycerol

Product overview

Description	Anti-HMGCL rabbit polyclonal antibody
Applications	ELISA, WB, IHC
Immunogen	Fusion protein of human HMGCL
Reactivity	Human, Mouse, Rat
Content	0.1 mg/ml
Host species	Rabbit
Ig class	Immunogen-specific rabbit IgG
Purification	Antigen affinity purification

Target information

Symbol	HMGCL
Full name	3-hydroxymethyl-3-methylglutaryl-CoA lyase
Synonyms	HL
Swissprot	P35914

Target Background

The protein encoded by this gene belongs to the HMG-CoA lyase family. It is a mitochondrial enzyme that catalyzes the final step of leucine degradation and plays a key role in ketone body formation. Mutations in this gene are associated with HMG-CoA lyase deficiency. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.

订购热线: 4008-898-798



Applications

Immunohistochemistry

Predicted cell location: Cytoplasm

Positive control: Human thyroid cancer

Recommended dilution: 50-200





The image on the left is immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using ml122488(HMGCL Antibody) at dilution 1/20, on the right is treated with fusion protein. (Original magnification: $\times 200$)

Predicted cell location: Cytoplasm

Positive control: Human prostate cancer

Recommended dilution: 50-200



The image on the left is immunohistochemistry of paraffin-embedded Human prostate cancer tissue using ml122488(HMGCL Antibody) at dilution 1/20, on the right is treated with fusion protein. (Original magnification: $\times 200$)

Western blotting

Predicted band size: 34 kDa

Positive control: SKOV3 cells, mouse heart and brain tissue

Recommended dilution: 200-1000

Gel: 8%SDS-PAGE


Lysate: 40 μ g

Lane 1-3: SKOV3 cells, mouse heart tissue, Mouse brain tissue

Primary antibody: ml122488(HMGCL Antibody) at dilution 1/150

Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution

Exposure time: 5 seconds



ELISA

Recommended dilution: 1000-2000

联系电话: 4008-898-798, 021-61725725

联系QQ: 2881505695, 2881505696

邮箱: mlbio_cn@yeah.net

网址: www.mlbio.cn